

Ignacio Zaballa,¹ Yeinzon Rodríguez,^{2,3} and David H. Lyth¹

¹*Department of Physics, Lancaster University, Lancaster LA1 4YB, United Kingdom*

²*Escuela de Física, Universidad Industrial de Santander, Ciudad Universitaria, Bucaramanga, Colombia*

³*Centro de Investigaciones, Universidad Antonio Nariño, Cll 58A # 37 - 94, Bogotá D.C., Colombia*

In this paper we calculate additional contributions to that part of the non-Gaussianity of the primordial curvature perturbation ζ , which come from the three-point correlator of the field perturbations. We estimate this contribution in the following models for the origin of ζ : single-component inflation, multi-component chaotic inflation, a two-component “hybrid” inflationary model, and the curvaton scenario. In all of these models, the additional contributions to the primordial non-gaussianity considered here are too small to ever be detected.

PACS numbers: 98.80.-k

JCAP 001P 0406, astro-ph/0603534

I. INTRODUCTION

There is substantial interest in an observable deviation from a Gaussian distribution of the primordial curvature perturbation ζ [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15]. From observation, the nongaussian contribution of ζ is expected to be small [16]. Nevertheless the current bound still includes a wide range of models for the origin of ζ , and therefore a future nongaussian detection or a tighter bound on the nongaussianity of ζ , could rule out a considerable number of models.

The level of nongaussianity in the curvature perturbation can be parameterised by f_{NL} , which specifies the three-point correlator of ζ . Here we apply the δN formalism [17, 18] to calculate new contributions to f_{NL} . While the relevant scales are outside the horizon, the curvature perturbation according to the δN formalism, is given by [5]

$$\zeta(\mathbf{x}, t) = \delta N(\rho(t), \phi_i(\mathbf{x})) \quad (1)$$

$$= \sum_i N_i(t) \delta \phi_i(\mathbf{x}) + \frac{1}{2} \sum_{ij} N_{ij}(t) \delta \phi_i(\mathbf{x}) \delta \phi_j(\mathbf{x}) + \dots, \quad (2)$$

where $N(\rho(t), \phi_i(\mathbf{x}))$ is the number of e-foldings from an initial flat slice, on which the field values are evaluated, to an uniform energy density slice at a time t after ζ has settled down to the time-independent value. N_i and N_{ij} are the partial derivatives, $N_i = \partial N / \partial \phi^i$ and $N_{ij} = \partial^2 N / \partial \phi^i \partial \phi^j$, resulting from the Taylor expansion on the field perturbations. From now on, we consider only the first two terms of the expansion in Eq.(2)

In the case of Gaussian field perturbations f_{NL} is given by [5, 6]

$$\frac{6}{5} f_{\text{NL}} \simeq \frac{\sum_{ij} N_i N_j N_{ij}}{(\sum_l N_l^2)^2} + \mathcal{P}_\zeta \frac{\sum_{ijk} N_{ij} N_{jk} N_{ki}}{(\sum_l N_l^2)^3}. \quad (3)$$

Seery and Lidsey [8] have calculated the three-point correlator of the field perturbations. To leading order in the field perturbations, the three-point correlator is

$$\langle \zeta_{\mathbf{k}_1} \zeta_{\mathbf{k}_2} \zeta_{\mathbf{k}_3} \rangle \subset \sum_{ijk} N_i N_j N_k \langle \delta \phi_{\mathbf{k}_1}^i \delta \phi_{\mathbf{k}_2}^j \delta \phi_{\mathbf{k}_3}^k \rangle, \quad (4)$$

which gives the following additional term

$$\Delta f_{\text{NL}} = \frac{\sum_{ijk} N_i N_j N_k f^{ijk}(k_1, k_2, k_3)}{[\sum_l N_l^2]^{3/2} \mathcal{P}_\zeta^{1/2}}. \quad (5)$$

The f^{ijk} 's are dimensionless functions proportional to the three-point correlator of the field perturbations. This term adds a contribution to f_{NL} too small to ever be observable, $|\Delta f_{\text{NL}}| \leq 0.044$ [11].

In this paper we calculate the next order contribution to f_{NL} . Contrary to the Δf_{NL} case, we are unable to give an estimate for an arbitrary potential. Therefore we calculate its contribution to the nongaussianity of ζ in several specific cases: single field inflation, multicomponent chaotic inflation, the two-component inflationary model of Ref. [19], and in the curvaton scenario.

The observed distribution of the curvature perturbation ζ in the CMB is almost Gaussian. The power spectrum in the δN formalism is then approximately given by

$$\mathcal{P}_\zeta \simeq \left(\frac{H}{2\pi} \right)^2 \sum_i N_i^2. \quad (6)$$

The three-point correlator of ζ , or its bispectrum defined by $\langle \zeta_{\mathbf{k}_1} \zeta_{\mathbf{k}_2} \zeta_{\mathbf{k}_3} \rangle = (2\pi)^3 B_\zeta \delta^3(\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3)$, is the lowest order signature of non-Gaussianity. To measure a possible small deviation from gaussianity in ζ we introduce the f_{NL} function, defined as [3]

$$B_\zeta(k_1, k_2, k_3) = \frac{6}{5} f_{NL} [P_\zeta(k_1) P_\zeta(k_2) + \text{cyclic}], \quad (7)$$

where $P_\zeta(k) = 2\pi^2 \mathcal{P}_\zeta / k^3$.

We consider the contribution of the three-point correlator of the field perturbations to the following term¹

$$\langle \zeta_{\mathbf{k}_1} \zeta_{\mathbf{k}_2} \zeta_{\mathbf{k}_3} \rangle \subset \frac{1}{4} \sum_{ijk} N_i N_j N_k \langle \delta\phi_{\mathbf{k}_1}^i (\delta\phi_{\mathbf{k}_2}^j)^2 (\delta\phi_{\mathbf{k}_3}^k)^2 \rangle + (k_1 \leftrightarrow k_2) + (k_1 \leftrightarrow k_3), \quad (8)$$

where $k_1 \leftrightarrow k_2$ and $k_1 \leftrightarrow k_3$ represent the corresponding permutations, and $(\delta\phi_{\mathbf{k}}^2)$ denotes the convolution product of the field perturbations:

$$(\delta\phi^2)_{\mathbf{k}} = \frac{1}{(2\pi)^3} \int d^3\mathbf{q} \delta\phi_{\mathbf{q}} \delta\phi_{\mathbf{k}-\mathbf{q}}. \quad (9)$$

The correlator in the field perturbations has the following generic form

$$\langle \delta\phi_{\mathbf{k}_1}^i \delta\phi_{\mathbf{q}_2}^j \delta\phi_{\mathbf{k}_2-\mathbf{q}_2}^j \delta\phi_{\mathbf{q}_3}^k \delta\phi_{\mathbf{k}_3-\mathbf{q}_3}^k \rangle = \langle \delta\phi_{\mathbf{k}_1}^i \delta\phi_{\mathbf{q}_2}^j \rangle \langle \delta\phi_{\mathbf{k}_2-\mathbf{q}_2}^j \delta\phi_{\mathbf{q}_3}^k \delta\phi_{\mathbf{k}_3-\mathbf{q}_3}^k \rangle + \text{Comb}, \quad (10)$$

where Comb refers to all possible combinations of the 5 elements in groups of two, which in total number 10. In the equation above, the two point function is defined by

$$\langle \delta\phi_{\mathbf{k}_1}^i \delta\phi_{\mathbf{k}_2}^j \rangle = (2\pi)^3 \delta^{ij} P_{\delta\phi^i}(k_1) \delta^{(3)}(\mathbf{k}_1 + \mathbf{k}_2), \quad (11)$$

where we consider a scale invariant spectrum $\mathcal{P}_{\delta\phi^i}(k) = (k^3/2\pi^2) P_{\delta\phi^i} = (H/2\pi)^2$. On the other hand, the three-point function obtained from second order cosmological perturbation theory is [8]

$$\langle \delta\phi_{\mathbf{k}_1}^i \delta\phi_{\mathbf{k}_2}^j \delta\phi_{\mathbf{k}_3}^k \rangle = (2\pi)^3 \frac{\delta^{(3)}(\sum_i \mathbf{k}_i)}{\prod_l k_l^3} \frac{4\pi^4}{M_P^2} \left(\frac{H}{2\pi} \right)^4 \sum_{\sigma'} \frac{\dot{\phi}_i}{2H} \delta_{jk} \mathcal{M}_{123}, \quad (12)$$

with $\mathcal{M}_{123}$ given by

$$\mathcal{M}_{123} \equiv \mathcal{M}(k_1, k_2, k_3) = \frac{1}{2} \left[-3 \frac{k_2^2 k_3^2}{k_t} - \frac{k_2^2 k_3^2}{k_t^2} (k_1 + 2k_3) + \frac{1}{2} k_1^3 - k_1 k_2^2 \right], \quad (13)$$

where k_t denotes the sum of the modulus of the three momenta $k_t = k_1 + k_2 + k_3$. In Eq. (12) the sum σ' is over all the permutations of the indices i, j and k , and at the same time over their respective momenta k_1, k_2 and k_3 . M_P is the reduced Plank mass: $M_P = 2.436 \times 10^{18}$ GeV.

Of the 10 combinations in Eq. (10) only 8 are non-vanishing, since neither $\mathbf{k}_2$ nor $\mathbf{k}_3$ can be zero. The 8 terms can be arranged into two groups of 4 terms, each group containing terms that give the same contribution. These can be represented schematically as two different diagrams, shown in Fig.(1). The other two contributions in Eq. (8) arising

¹ Note that we have diagonalized the N_{ij} in the field perturbations basis to simplify the calculation. The connected part of the five point function is not written explicitly.




FIG. 1: Here we show the two possible diagrams. There are three external lines corresponding to the three momenta k_1 , k_2 and k_3 . The parallel lines represent the two-point correlator, whereas the crossed circle indicates the three-point correlator. The diagrams have been drawn using JaxoDraw [20].

from permuting the k_1 momentum with k_2 and k_3 , result in the same two diagrams. Similar diagrams are given in Ref. [21].

In calculating the contribution to the three-point function of ζ resulting from the first diagram, we find the following integral over the internal momentum:

$$I_{123} = \int_{L^{-1}} \frac{d^3 \mathbf{q}}{q^3} \frac{\mathcal{M}(k_1, |\mathbf{k}_2 + \mathbf{q}|, |\mathbf{k}_3 - \mathbf{q}|)}{k_1^3 |\mathbf{k}_2 + \mathbf{q}|^3 |\mathbf{k}_3 - \mathbf{q}|^3}, \quad (14)$$

and similar integrals corresponding to the permutations in the argument of the $\mathcal{M}_{123}$ function. The momenta dependence of the integrand can be read off from the location of the three-point correlator in Fig.(1). In the equation above L^{-1} indicates that the integrand is set equal to zero in a sphere of radius L^{-1} around each singularity. The integral then is approximately given at each singularity by

$$I_{123} = 4\pi \ln KL \left(\frac{\mathcal{M}(k_1, |\mathbf{k}_2 + \mathbf{q}|, |\mathbf{k}_3 - \mathbf{q}|)}{k_1^3 |\mathbf{k}_2 + \mathbf{q}|^3 |\mathbf{k}_3 - \mathbf{q}|^3} \right)_{\mathbf{q}=0, -\mathbf{k}_2 \text{ or } \mathbf{k}_3}, \quad (15)$$

where the function in brackets is evaluated at the singularity, and $K = \min\{k_i\}$. The logarithm in the expression above can be taken to be of order 1 [6]. It can be shown that each permutation of the $\mathcal{M}_{123}$ function only diverges at $q = 0$, and only at one of the other two singularities if at any, since the integrand becomes 0 in at least one of them. Therefore we are able to give a momenta dependent result for each integral.

The total contribution to the f_{NL} function from the first diagram, using the identity $\sum_i N_i \dot{\phi}^i = -H$ as in [15], gives

$$\frac{6}{5} \Delta f_{NL}^{(1)} = \frac{1}{M_p^2} \left(\frac{H}{2\pi} \right)^4 \frac{\left[2 \sum_i N_i N_{ii}^2 \left(\frac{\dot{\phi}^i}{2H} \right) - \frac{1}{2} \sum_j N_{jj}^2 \right] \frac{\sum_\sigma \mathcal{M}_{123}}{\sum_l k_l^3} - 4 \sum_i N_i N_{ii}^2 \left(\frac{\dot{\phi}^i}{2H} \right)}{\sum_k N_k^2}. \quad (16)$$

The momenta dependent term has a maximum absolute value for $k \equiv k_1 = k_2 = k_3$ given by [11]

$$\frac{\sum_\sigma \mathcal{M}(k, k, k)}{\sum_l k_l^3} = -\frac{11}{6}, \quad (17)$$

where the sum over σ denotes the sum over the permutations of the three momenta. In the following sections, we calculate the value of $\Delta f_{NL}^{(1)}$ for the maximum of the momenta dependent term.

The integrals for the second diagram are evaluated in a similar way to the integrals of the first diagram, but this time there are more cancellations at the singularities. The total contribution to f_{NL} function from the second diagram is

$$\frac{6}{5} \Delta f_{NL}^{(2)} = -\frac{4}{M_p^2} \left(\frac{H}{2\pi} \right)^4 \left[\frac{\sum_i N_i N_{ii}^2 \left(\frac{\dot{\phi}^i}{2H} \right)}{\sum_k N_k^2} \right]. \quad (18)$$

In the following sections, we evaluate these higher order contributions in Eqs. (16) and (18) for several specific cases.

In single field inflation the number of e-foldings $N(t, t_0)$ is given by

$$N(t, t_0) \equiv \int_{t_0}^t H dt \simeq \frac{1}{M_p^2} \int_{\phi}^{\phi_0} \frac{V}{V_\phi} d\phi. \quad (19)$$

In the δN formalism we evaluate the partial derivatives of N respect of the field values on the initial flat slice. Then

$$N_\phi = \frac{1}{\sqrt{2\epsilon} M_p}, \quad (20)$$

$$N_{\phi\phi} = \frac{1}{M_p^2} \left(1 - \frac{\eta}{2\epsilon} \right), \quad (21)$$

where $\epsilon = M_p^2/2 (V_\phi/V)^2$ and $\eta = M_p^2(V_{\phi\phi}/V)$ are the slow-roll parameters.

Defining $\Delta f_{NL}^{(1+2)} \equiv \Delta f_{NL}^{(1)} + \Delta f_{NL}^{(2)}$, we have the following contribution to f_{NL} :

$$\frac{6}{5} \Delta f_{NL}^{(1+2)} = 54 \mathcal{P}_\zeta^2 \left(\epsilon^3 - \eta\epsilon^2 + \frac{\eta^2\epsilon}{4} \right). \quad (22)$$

This is much less than the leading order contribution calculated in [8].

IV. MULTI-COMPONENT CHAOTIC INFLATION

In this section we consider a potential with the following form [13, 22]

$$V = \frac{1}{2} \sum_{i=1}^n m_i^2 \phi_i^2. \quad (23)$$

This potential represents multi-component chaotic inflation. Under slow-roll conditions the masses of the fields are similar [23], and then the number of e-foldings N is given by $N(\phi_i, \rho) = \frac{1}{4} \sum_{i=1}^n (\phi_i^2/M_P^2)$ [24]. The derivatives of N are

$$N_i = \phi_i/2M_p^2, \quad (24)$$

$$N_{ij} = \delta_{ij}/2M_p^2. \quad (25)$$

Then the contribution in Eq. (3) to f_{NL} is much less than one [13]; $(6/5)f_{NL} = 1/N + O(1/N^3)$, and therefore it is not observable. The higher order contributions in Eqs. (16) and (18) give

$$\frac{6}{5} \Delta f_{NL}^{(1+2)} = \left(\frac{r}{16} \right)^3 \mathcal{P}_\zeta^2 \left[\frac{11}{6} (n+2) + 8 \right], \quad (26)$$

where r is the tensor to scalar ratio $r = 8(H/2\pi)^2/M_p^2 \mathcal{P}_\zeta$, which in this case is given by $r = 8/N$. This contribution is much smaller than the leading order one in Eq. (3), and therefore negligible.

V. TWO-COMPONENT MODEL

Here we consider a two-component model given by the following potential [4, 19]:

$$V(N, \phi, \sigma) = V_0 \left[1 + \frac{1}{2} \eta_\phi \frac{\phi^2(N)}{M_p^2} + \frac{1}{2} \eta_\sigma \frac{\sigma^2(N)}{M_p^2} \right], \quad (27)$$

where η_ϕ and η_σ are the η slow-roll parameters. $\phi(N)$ and $\sigma(N)$ are given in terms of the field values just after horizon exit, ϕ and σ , by $\phi(N) = \phi \exp(-N\eta_\phi)$ and $\sigma(N) = \sigma \exp(-N\eta_\sigma)$. Here we consider the case $\sigma = 0$ of Ref. [19]. The derivatives of N are [5, 25]

$$N_\phi = (1/\eta_\phi \phi), \quad N_\sigma = 0, \quad (28)$$

$$N_{\phi\phi} = -\eta_\phi (1/\eta_\phi \phi)^2, \quad (29)$$

$$N_{\sigma\sigma} = \eta_\sigma (1/\eta_\phi \phi)^2 e^{2N(\eta_\phi - \eta_\sigma)}. \quad (30)$$

The f_{NL} function is dominated by the term in Eq. (3) proportional to $N_{\sigma\sigma}$, and it is approximately given by

$$\frac{6}{5}f_{NL} = \left(\frac{r}{16}\right)^3 \mathcal{P}_\zeta \left[\left(\frac{\eta_\sigma}{\epsilon}\right)^3 e^{6N(\eta_\phi - \eta_\sigma)} \right], \quad (31)$$

where ϵ is the slow roll parameter across the $\sigma = 0$ trajectory, $\epsilon = (\eta_\phi \phi / \sqrt{2} M_p)^2$, and r is the tensor to scalar ratio defined earlier. Similarly, $\Delta f_{NL}^{(1+2)}$ is dominated by the term proportional to $N_{\sigma\sigma}$, and therefore the main contribution comes from $\Delta f_{NL}^{(1)}$:

$$\frac{6}{5}\Delta f_{NL}^{(1)} = \frac{11}{6} \left(\frac{r}{16}\right)^3 \mathcal{P}_\zeta^2 \left[\left(\frac{\eta_\sigma}{\epsilon}\right)^2 e^{4N(\eta_\phi - \eta_\sigma)} \right]. \quad (32)$$

This contribution is much smaller than f_{NL} ;

$$\frac{\Delta f_{NL}^{(1)}}{f_{NL}} \simeq \mathcal{P}_\zeta \left(\frac{\eta_\sigma}{\epsilon}\right)^{-1} e^{-2N(\eta_\phi - \eta_\sigma)}. \quad (33)$$

VI. CURVATON SCENARIO

In the curvaton scenario, the curvature perturbation ζ is generated during the oscillations of the curvaton field σ after the end of inflation [26, 27, 28]. The curvaton starts oscillating around the minimum of a quadratic potential, $V(t, \mathbf{x}) = m^2 \sigma^2(t, \mathbf{x})/2$, in an initially radiation dominated epoch with amplitude $g \equiv g(\sigma_*)$ until it eventually decays. Here σ_* is the value of the curvaton field just after horizon exit during inflation. Defining N as the number of e-foldings from the beginning of the oscillations to the curvaton decay, its derivatives are given by [5]

$$N_{\sigma_*} = \frac{2\Omega_\sigma g'}{3g}, \quad (34)$$

$$N_{\sigma_*\sigma_*} = \frac{2\Omega_\sigma}{9g^2} [2\Omega_\sigma (2 + \Omega_\sigma) g'^2 - 3(g'^2 + gg'')], \quad (35)$$

where g' and g'' denote the first and second derivatives with respect to σ_* , and Ω_σ is the final fraction of the curvaton energy density before it decays. Then [4, 5]

$$f_{NL} = -\frac{5}{3} - \frac{5}{6}\Omega_\sigma + \frac{5}{4}\Omega_\sigma \left(1 + \frac{gg''}{g'^2}\right). \quad (36)$$

Defining $\epsilon_\sigma = M_p^2 (V_\sigma / \sqrt{2}V)^2$, where $V_\sigma = \partial V / \partial \sigma$, the relative contribution of $\Delta f_{NL}^{(1+2)}$ to f_{NL} is

$$\left(\frac{\Delta f_{NL}^{(1+2)}}{f_{NL}}\right) \simeq r \mathcal{P}_\zeta^2 f_{NL}, \quad (37)$$

being too small to be detected because of the current observational constraint, $f_{NL} < 121$ [16].

VII. CONCLUSION

We have calculated a higher-order contribution of the three-point correlator of the field perturbations to the three-point correlator of the curvature perturbation, which is the next to leading order in the field perturbations. Even though intuition suggests that it should be smaller than the leading-order contribution, we have not been able to give a general proof that this is so. Instead, we have shown that it is negligible for the cases that have been considered so far in the literature. Keeping only the first two terms of Eq.(2), there is one higher order contribution, and more contributions will be generated by higher order terms in Eq.(2). All of these contributions can be calculated in a similar way to the calculation shown here.

IZ is partially supported by Lancaster University Physics Department. YR is supported by COLCIEN-CIAS (COLOMBIA). DHL is supported by PPARC grants PPA/G/S/2002/00098, PPA/G/O/2002/00469, PPA/Y/S/2002/00272, PPA/V/S/2003/00104 and EU grant MRTN-CT-2004-503369.

-
- [1] N. Bartolo, S. Matarrese, and A. Riotto, Phys. Rev. D **65**, 103505 (2002) [arXiv:hep-ph/0112261].
 - [2] V. Acquaviva, N. Bartolo, S. Matarrese, and A. Riotto, Nucl. Phys. B **667**, 119 (2003) [arXiv:astro-ph/0209156].
 - [3] J. Maldacena, JHEP **0305**, 013 (2003) [arXiv:astro-ph/0210603].
 - [4] D. H. Lyth and Y. Rodriguez, Phys. Rev. D **71**, 123508 (2005) [arXiv:astro-ph/0502578].
 - [5] D. H. Lyth and Y. Rodriguez, Phys. Rev. Lett. **95**, 121302 (2005) [arXiv:astro-ph/0504045].
 - [6] L. Boubekeur and D. H. Lyth, Phys. Rev. D **73**, 021301(R) (2006) [arXiv:astro-ph/0504046].
 - [7] Y. Rodríguez, PhD Thesis, Lancaster University, Lancaster UK (2005) [arXiv:astro-ph/0507701].
 - [8] D. Seery and J. E. Lidsey, JCAP **0509**, 011 (2005) [arXiv:astro-ph/0506056].
 - [9] G. I. Rigopoulos, E. P. S. Shellard, and B. J. W. van Tent, Phys. Rev. D **72**, 083507 (2005) [arXiv:astro-ph/0410486].
 - [10] G. I. Rigopoulos, E. P. S. Shellard, and B. J. W. van Tent, [arXiv:astro-ph/0506704, arXiv:astro-ph/0511041].
 - [11] D. H. Lyth and I. Zaballa, JCAP **0510**, 005 (2005) [arXiv:astro-ph/0507608].
 - [12] L. E. Allen, S. Gupta, and D. Wands, JCAP **0601**, 006 (2006) [arXiv:astro-ph/0509719].
 - [13] L. Alabidi and D. H. Lyth, [arXiv:astro-ph/0510441].
 - [14] N. Barnaby and J. M. Cline, [arXiv:astro-ph/0601481].
 - [15] F. Vernizzi and D. Wands, [arXiv:astro-ph/0603799].
 - [16] E. Komatsu *et al.*, Astrophys. J. Suppl. Ser. **148**, 119 (2003) [arXiv:astro-ph/0302223].
 - [17] M. Sasaki and E. D. Stewart, Prog. Theor. Phys. **95**, 71 (1996) [arXiv:astro-ph/9507001].
 - [18] D. H. Lyth, K. A. Malik, and M. Sasaki, JCAP **0505**, 004 (2005) [arXiv:astro-ph/0411220].
 - [19] K. Enqvist and A. Vähikönen, JCAP **0409**, 006 (2004) [arXiv:hep-ph/0405103].
 - [20] D. Binosi and L. Theussl, Comput. Phys. Commun. **161**, 76 (2004) [arXiv:hep-ph/0309015].
 - [21] M. Crocce and R. Scoccimarro, Phys. Rev. D **73**, 063519 (2006) [arXiv:astro-ph/0509418].
 - [22] S. Dimopoulos, S. Kachru, J. McGreevy and J. G. Wacker, [arXiv:hep-th/0507205].
 - [23] D. Polarski and A. A. Starobinsky, Nucl. Phys. B **385**, 623 (1992).
 - [24] D. H. Lyth and A. Riotto, Phys. Rep. **314**, 1 (1999) [arXiv:hep-ph/9807278].
 - [25] K. A. Malik, JCAP **0511**, 005 (2005) [arXiv:astro-ph/0506532].
 - [26] D. H. Lyth and D. Wands, Phys. Lett. B **524**, 5 (2002) [arXiv:hep-ph/0110002].
 - [27] T. Moroi and T. Takahashi, Phys. Lett. B **522**, 215 (2001) [Erratum-ibid. B **539**, 303 (2002)] [arXiv:hep-ph/0110096].
 - [28] D. H. Lyth, C. Ungarelli, and D. Wands, Phys. Rev. D **67**, 023503 (2003) [arXiv:astro-ph/0208055].